

SOME POST HARVEST INSECT PESTS THAT THREATEN FOOD SECURITY

Seed Enterprises Management Institute

University of Nairobi
Prof. Florence Olubayo



Introduction

- Factors such as storage duration, prevailing environmental conditions and crop varieties influence insect populations development and losses incurred.
- Inadequate storage methods lead to losses in stored grain sometimes of unacceptable magnitude in SSA.
- These pests inflict both direct and indirect damage to the grain, and the most important ones start in the field.

Damage caused

Direct damage

- Kernel damage,
- Contamination,
- Grain dust,
- Damage to wooden structures and other containers

Indirect damage

- Dry grain heating and moisture migration in storage
- Lowered germination of seed grains
- Distribution of molds and other organisms through the grain mass
- Insect fragments in cereal products

THE GRAIN MOTH

(Sitotroga cerealella (ol.))

- Small straw coloured moth (wing span 10-18mm)
- Able to fly from infested grain in store to the field
- Infests maturing cereals in the field
- Infestation can also occur at store levels.
- Damage:- small circular 'windows' and holes on the grain
- Causes severe damage to cereals stored mainly in unthreshed form
- Grain attacked:- maize, sorghum, wheat, paddy & barley.

SITOTROGA CEREALELLA
(Angoumois Grain Moth)



MAIZE AND RICE WEEVILS

(Sitophilus spp.)

- Dark brown weevils (2.5-4.5mm long)
- Able to fly from infested grain in store to the field.
- Infests maturing cereals in the field
- Infestation can also occur at store level
- Damage: Small circular holes on the surface of the grain
- Causes severe damage to grain stored in both threshed and unthreshed form
- Grain attacked:- maize, millet, sorghum, wheat, barley and rice




***SITOPHILUS* spp.**
(Maize and Rice Weevils)

LARGER GRAIN BORER

(Prostephanus truncatus(h.))

- A dark brown cyclindrical beetle (3-4.5mm long)
- Able to fly from infested grain in store to the field
- Infests maize in the field before harvest
- Infestation can also occur at store level
- The beetle eats funnels and holes in the husks, grain and cob.
- Very serious pest that also eats into the wooden store structures
- Also feeds on dried cassava



PROSTEPHANUS TRUNCATUS
(Larger Grain Borer)

PULSE BRUCHIDS (BEETLES)

- THE BEAN BRUCHID (*Acanthoscelides obtectus* (say)).
- Grey to brown oval beetles (3 – 4.5mm long)
- Able to fly from infested grain the the store to the field
- Infestation can also occur at store level
- Damage:- small dark ‘windows’ and holes on the grain
- Causes serious damage to stored beans

ACANTHOSCELIDES OBTECTUS

(Bean Beetle)



THE COWPEA BRUCHIDS

(Callosobruchus spp)

- Light to dark brown beetles (2.0-3.5mm long)
- Able to fly from infested grain in stores to the field
- Infests maturing legumes (cowpea, pigeon peas, chick peas and grains) in the field
- Infestation can also occur at store level
- Small dark ‘windows’ and holes on the grain indicate infestation by the bruchids
- Causes serious damage to stored pulses.

Callosobruchus spp.



THE FLOUR BEETLE


(Tribolium spp)

- Reddish brown flat beetles (2.5 – 4.5mm long)
- Infests stored (broken) grain and milled products
- Causes high level of gram contamination
- Presence of reddish brown beetles, cast skins and faecal pellets on damaged grain and milled products indicates infestation by these beetles.
- Serious secondary pests of all stored grain and milled grain products



TRIBOLIUM CASTANEUM
(Rust-Red Flour Beetle)

SEMISUON
Seed Enterprises Management Institute
University of Nairobi



SEM JON

Seed Enterprises Management Institute

EPHESTIA spp.
(Tropical Warehouse Moths)

Several species of *Ephestia* may be encountered in tropical stores. They attack a wide range of products particularly damaged or processed cereals, dried fruit, nuts, cocoa and even tobacco. Only the larvae feed. They also leave trails of silk which can form a thick webbing over and in the stored food. Reconditioning food to remove webbing can be very costly.

(Wing span 11-28 mm)

Some Management Practices

Post-harvest Insect pest control should begin before the crop is mature and must definitely begin before it is harvested and put in drying structures.

Proper program for insect control include:

- Select plant varieties with good husk cover and inherent resistance to field and storage pests

- Repair the store and thoroughly clean before the new crop is mature.
- Clear the surroundings of the store of any waste that can harbor insect pests
- Harvest early to avoid field infestation
- Dry the grain as fast as possible and shell it when dry
- Shell carefully to avoid damage to the kernels
- Treat the dry grain with an appropriate insecticide

- Carry out regular inspections of the stored grain to detect any infestation and take control measures as necessary
- Carry out principles of good store management, including maintenance, stock rotation and hygiene.

Seed Enterprises Management Institute

University of Nairobi