

**NUTRITIONAL KNOWLEDGE AND DIETARY PRACTICES  
OF PREGNANT WOMEN RECEIVING NUTRITION EDUCATION  
WHILE ATTENDING ANTENATAL CARE AT MBAGATHI HOSPITAL  
NAIROBI COUNTY KENYA**

**MARYGORRET NELIMA MALOBA**

**A DISSERTATION SUBMITTED IN PARTIAL FULFILMENT OF THE  
REQUIREMENTS FOR THE AWARD OF THE DEGREE OF MASTER  
OF SCIENCE IN APPLIED HUMAN NUTRITION**

**DEPARTMENT OF FOOD SCIENCE NUTRITION AND TECHNOLOGY  
FACULTY OF AGRICULTURE  
UNIVERSITY OF NAIROBI**

**2022**

## DECLARATION

This dissertation is my original work and has not been submitted forward of degree in any other University.

Signature;

Date: 28.08.2022




Signature 

Date: 28.08.2022.

**Professor Wambui Kogi-Makau**

Department of Food Science Nutrition and Technology

University of Nairobi

Signature   
í í í í í í í í .

Date: 28.08. 2022

**Dr Dasel Wambua Mulwa Kaindi**

Department of Food Science Nutrition and Technology

University of Nairobi



**UNIVERSITY OF NAIROBI**

**PLAGIARISM DECLARATION FORM**

**Name of Student:** MARYGORRET NELIMA MALOBA

**Registration Number:** A56/6750/2017

**College:** AGRICULTURE AND VETERINARY SCIENCES

**Faculty:** AGRICULTURE

**Department:** FOOD SCIENCE NUTRITION AND TECHNOLOGY

**Course Name:** MASTER OF SCIENCE IN APPLIED HUMAN NUTRITION

**Title of the Work:** NUTRITIONAL KNOWLEDGE AND DIETARY PRACTICES  
OF PREGNANT WOMEN RECEIVING NUTRITION EDUCATION WHILE  
ATTENDING ANTENATAL CARE AT MBAGATHI HOSPITAL NAIROBI  
COUNTY

**DECLARATION**

1. I understand what Plagiarism is and I am aware of the University's policy in this regard
2. I declare that this dissertation is my original work and has not been submitted elsewhere for examination, award of a degree or publication. Where other people's work, or my own work has been used, this has properly been acknowledged and referenced in accordance with the University of Nairobi requirements.
3. I have not sought or used the services of any professional agencies to produce this work
4. I have not allowed, and shall not allow anyone to copy my work with the intention of passing it off as his/her own work
5. I understand that any false claim in respect of this work shall result in disciplinary action, in accordance with University Plagiarism Policy.

Signature;

Date: 28.08.2022

## TABLE OF CONTENTS

DECLARATION .....	ii
LIST OF TABLES .....	vi
LIST OFFIGURES .....	vii
ABSTRACT.....	viii
ACRONYMS AND ABBREVIATIONS .....	x
OPERATIONAL DEFINITION .....	xi
CHAPTER ONE: INTRODUCTION.....	1
1.1 Background Information .....	1
1.2 Statement of the Problem .....	3
1.3 Justification of the study .....	3
1.4 Aim of the study .....	3
1.5 Purpose of the study.....	3
1.6 Objectives of the study.....	4
1.6.1 Main Objective .....	4
1.6.2 Specific objectives .....	4
1.7 Research Questions.....	4
CHAPTER TWO: LITERATURE REVIEW .....	5
2.0 Introduction .....	5
2.1 Overview of Antenatal Care .....	5
2.2 Nutritional knowledge of pregnant women .....	7
2.3 Dietary intake of pregnant women.....	8
2.4 Research Design and methodology.....	11
2.4 Gaps in knowledge.....	13
2.6 Conceptual Framework of the study variables .....	15
CHAPTER THREE: RESEARCH DESIGN AND METHODOLOGY .....	16
3.1 Study Design .....	16
3.2 Study site and Population.....	16
3.2.1 Study site .....	16
3.2.2 Study population .....	17
3.3 Sample size.....	17
3.3.1 Sample size determination.....	17
3.3.2 Sampling procedure .....	18
3.3.3 Inclusion criteria .....	18
3.3.4 Exclusion criteria .....	19
3.4 Data collection tools and procedure.....	19

3.4.1 Study Questionnaire.....	19
3.5 Recruitment and Training of Research Assistants .....	21
3.6 Pre -Test of the Data Collection Tools.....	21
3.7 Ethical and Human Rights Consideration .....	21
3.8 Data analysis and Quality Assurance Procedures .....	22
CHAPTER: FOUR RESULTS .....	23
4.1 Socio-demographic characteristics of the respondents .....	23
4.2 Gestational age and ANC attendance of pregnant women at Mbagathi hospital.....	24
4.3 Medical profile for the current pregnancy of pregnant women at Mbagathi hospital.....	25
4.4 Nutrition education topics taught to pregnant women attending antenatal care at Mbagathi Hospital.....	26
4.5 Level of nutrition Knowledge of pregnant women attending antenatal care at Mbagathi hospital .....	27
4.6 Dietary Diversity score of pregnant women attending antenatal care at Mbagathi hospital .....	28
4.7 Nutritional status of the pregnant women attending antenatal care at Mbagathi hospital.....	32
CHAPTER FIVE: DISCUSSION .....	34
5.1 Introduction .....	34
5.2 Socio-demographic context of the study participants .....	34
5.3 Level of Nutritional knowledge of the study participants .....	35
5.4 Dietary practices of the pregnant women attending antenatal care at Mbagathi hospital .....	36
5.5 Distribution of nutritional knowledge and mean dietary diversity score of pregnant women attending antenatal care at Mbagathi hospital .....	36
5.6 Nutrition status of the study participants attending antenatal care at Mbagathi hospital .....	37
5.7 Distribution of nutritional knowledge levels by nutritional status of pregnant women attending antenatal care at Mbagathi hospital .....	37
5.8 Association between the level of nutritional knowledge and antenatal care attendance.....	38
CHAPTER SIX: CONCLUSIONS AND RECOMMENDATIONS .....	39
6.1 Conclusion.....	39
6.2 Recommendations .....	39
REFERENCE.....	40
APPENDIX 1: STUDY QUESTIONNAIRE .....	44
APPENDIX 2: OBSERVATION GUIDE AT THE ANTE NATAL CLINIC .....	48
APPENDIX 3: GUIDELINES AND RESOURCE MATERIALS AVAILABLE ON MATERNAL NUTRITION.....	49
APPENDIX 4: CONSENT FORM .....	50

## LIST OF TABLES

Table 1: Demographic characteristics of the respondents attending antenatal care at Mbagathi Hospital .....	23
Table 2: Gestational age and ANC attendance of pregnant women attending ANC at Mbagathi hospital.....	25
Table 3: Medical profile of the respondents of pregnant women .....	25
Table 4: Topics taught during nutrition education sessions at Mbagathi hospital .....	26
Table 5: Association between the level of nutritional knowledge and nutrition status (MUAC) .....	28
Table 6: Distribution of socio-demographic characteristics by dietary diversity score.....	30
Table 7: Association between nutritional knowledge and consumption of the recommended ten food groups and the Mean Dietary Diversity score of pregnant women attending antenatal care at Mbagathi hospital .....	31

## LIST OFFIGURES

Figure 1: Illustration of research study designs .....	<b>Error! Bookmark not defined.</b>
Figure 2: Knowledge-Attitude-Behavior (KAB) model .....	14
Figure 3: Conceptual framework derived from study objectives .....	15
Figure 4: Map of Kenya showing Mbagathi Hospital.....	17
Figure 5: Sampling frame for study site and study population at Mbagathi hospital .....	18
Figure 6: Level of nutritional knowledge of the study participants at Mbagathi hospital .....	28
Figure 7: Dietary Diversity Scores of the pregnant women attending antenatal care at Mbagathi hospital.....	29
Figure 8: Nutritional status of pregnant women attending antenatal care at Mbagathi hospital based on MUAC cut-off points.....	32
Figure 9: Association between the level of nutritional knowledge and nutritional status of the women attending antenatal care at Mbagathi hospital .....	33

## **ABSTRACT**

Nutrition education is one of the important components of antenatal care. It is a widely used strategy to improve nutritional knowledge of pregnant women seeking ante natal care more so in government run health care facilities. A lot of studies to assess the level of nutritional knowledge on pregnant women attending antenatal care in government health facilities have been done but little data exists on the antenatal care situation at Mbagathi hospital given that it is one of the key government run level five hospital facility in Nairobi County that serves a population of about 3.1 million urban dwellers. The broad objective of this study was to determine the nutritional knowledge and dietary practices of pregnant women receiving nutrition education while attending antenatal care at Mbagathi hospital Nairobi county.

The study used a descriptive cross-sectional study design for collection and analysis of data and. Total of 195 pregnant women receiving nutrition education while attending antenatal care in Mbagathi hospital Antenatal clinic were recruited into the study through systematic sampling. The study, using a semi structured questionnaire, assessed the level of nutritional knowledge and dietary practices of attendees of ante natal clinic. Anthropometric measurement of the Middle-upper arm Circumference (MUAC) was used to assess the nutrition status of the women. The study used Statistical Package for the Social Sciences (SPSS) version 20 software for data entry and analysis. Descriptive statistics (mean, median, frequency and standard deviation) described the background characteristics of the study sample while inferential statistics, (P-Value of <0.05 at 95% confidence interval), were employed to determine associations and relationships between two or more variables.

Majority (69%) of the participants were within the age category of between 21- 30 years and the mean age was 27 ( $\pm=5.38$ ) years. Over three quarters (79%) were married while 21% were not. Slightly below half of the women, 43%, had attained secondary education 17% had primary education while 9% had University education. Slightly over a third (35%) were housewives, 31% self-employed and 12% were salaried workers. The mean household size was 3.18 ( $\pm=1.47$ )

Over 50% of the respondents were in their third trimester of pregnancy, 34% in the second and 7% in the first trimester. Only 39% had made between three and four antenatal visits, with the mean number of visits was 3.7. Nearly all (94%) participants had blood pressure readings taken, 71% had a concurrent illness (colds and flus, urinary tract infections,



Malaria), 8% had a diagnosis of High Blood Pressure, 4% had gestational diabetes while 2% had iron deficiency anemia. On assessment of nutritional knowledge, 84% displayed adequate levels of nutrition knowledge with regards to aspects of nutrition during pregnancy like variety of foods; amount of food during pregnancy; energy sources; micronutrient supplement for pregnant women, necessity for supplementation and duration; micronutrients, iron and calcium and consequences of deficiency during pregnancy while 13% had poor knowledge score. The level of nutritional knowledge was strongly associated with the number of antenatal care visits attended ( $p=0.03$ ), nutrition status ( $p=0.018$ ) and the mean dietary diversity score ( $p=0.03$ ) of the participants. Assessment of dietary diversity showed that having consumed at least five food groups, 63% met the minimum dietary diversity score for women. There was a significant association between household size and dietary diversity score ( $p=0.033$ ). Majority (65%) of pregnant women from household sizes of 1 to 4 members displayed adequate dietary diversity. However, there was no association between dietary diversity with education status, marital status, occupation and level of income. Nutrition status was assessed by taking the reading of the middle upper arm circumference (MUAC) using an adult MUAC tape. The respondents had MUAC measurement of ranging from 22.8cm to over 30cm, with 62% of the pregnant women had normal nutritional status with a MUAC reading of between 22.8-30.7cm, 27% of the women were obese with a MUAC readings of >30cm while 11% with MUAC reading of <22 cm were underweight.

From the research findings, pregnant women who receive nutrition education while attending antenatal care ante natal care at Level 5 hospitals in Kenya have adequate maternal nutrition knowledge and dietary diversity score. The more the number of ante natal visits a pregnant woman makes, the higher the level of nutritional knowledge.

This study recommends that the Government of Kenya, Ministry of Health and partners should continuously and consistently create more awareness of the existence of antenatal care service and its benefits to pregnant women. Health workers at antenatal clinics should also pay particular attention to underweight and overweight pregnant women since they have poor nutrition knowledge the health workers should go an extra mile to find out why they are unable to make the right food choices

## **ACRONYMS AND ABBREVIATIONS**

ANC	Ante natal care
DDS	Dietary Diversity Score
DDS-W	Dietary Diversity Score for Women
FAO	Food and Agriculture Organization
FGD	Focus Group Discussion
IFAS	Iron and Folic acid supplements
KAB	Knowledge Attitude Behavior model
KDHS	Kenya Demographic and Health Survey
KII	KeyInformant Interview
MDD	Minimum Dietary Diversity
MDDS	Mean Dietary Diversity Score
MIYCN	Maternal Infant and Young Child Nutrition
MOH	Ministry of Health
MUAC	Mid-Upper Arm Circumference
WHO	World Health Organization
WRA	Women of Reproductive Age

## **OPERATIONAL DEFINITION**

**Antenatal care**-It is a form of health care where pregnant women learn from skilled health personnel about proper nutrition during pregnancy, they receive micronutrient supplements and treatment in case of an illness and also learn about healthy behavior during pregnancy. The health care personnel also monitor pregnancy and any danger signs during ante natal visits

**Dietary practices**-Dietary practice is defined as a participant's preference in food consumption or behavior of dietary habits and can be classified as good dietary practice and poor dietary practices

**Nutrition Education**-This is education given to pregnant women during antenatal care visits about their nutritional needs and requirements in pregnancy. Nutrition education covers variety of topics on maternal nutrition

**Nutritional knowledge**-Nutritional knowledge refers a pregnant woman understanding on various aspect of maternal nutrition including energy and protein requirements in pregnancy, micronutrient requirements, serving size of food groups, diversity of diets. The purpose of equipping pregnant women with nutritional knowledge is to help them improve their dietary practices in order to meet her nutritional needs together with the needs of the developing baby

**Nutritional Status**-This is the state of a pregnant woman's health that is determined by wrapping a tape otherwise known as an adult MUAC on the left Middle Upper Arm and reading the circumference in centimeters. The reading is categorized as either underweight (a reading of less than 22cm, overweight (a reading of over 30cm or normal (a reading of between 22-30cm)

**Middle Upper Arm Circumference**- This is a reading taken on the upper left arm of a pregnant woman in order to determine her nutrition status. The reading is taken in centimeters and categorized as either underweight, normal or overweight

# CHAPTER ONE: INTRODUCTION

## 1.1 Background Information

Globally, the prevalence of malnutrition in women, in all its forms, is still unacceptably high. Malnutrition (underweight, overweight, anemia and short stature) affects millions of women around the globe including during pregnancy and breastfeeding which are periods that are marked by increased nutritional needs

Globally, about 170 Million women (9.1%) are underweight and thrice as many, 610 Million (32.5%) are overweight. Prevalence of Malnutrition among women in Sub-Saharan and South Asia is still very high (UNICEF, 2022)

Ante natal Care (ANC) serves as a major entry point where pregnant women receive education for health promotion and prevention. In 2003, WHO recommended that pregnant women should attend a minimum of four ANC appointments throughout pregnancy, this was termed Focused Antenatal Care ideally at 16, 24-28, 32 and 36 weeks of pregnancy. One of the important components of ANC is health promotion including nutrition education among others (World Health Organization, 2003).

When women attend several ANC visits, they exhibit better nutritional knowledge, attitude and practices compared to those not attending more so in developing countries (Girard & Olude, 2012). Nutrition education is a widely used strategy to improve nutrition status of pregnant women.

One of the major areas that nutrition education focuses on is equipping pregnant women with knowledge that will enable them improve on the quality of their diet, as well as education on food types and amount of servings they need to consume to optimize dietary intake. Deficiencies in micronutrients e.g.

Iron, iodine, calcium, vitamin A and iodine can lead to pregnancy complications which put the woman and baby at risk. When pregnant women do not gain adequate weight because of insufficiency in the diet, they stand a risk of pregnancy outcomes like preterm deliveries, babies with a low birth weight and also getting babies with congenital malformations. Apart from education on nutritional requirements in pregnancy, pregnant women are also counselled on dangers of some lifestyle behaviors that may harm the unborn baby such as smoking, alcohol drinking and use of illegal drugs. Women attending ANC also benefit from a nutrition intervention of Iron and folic acid supplements (IFAS) as a preventive measure against Iron Deficiency Anemia and neural tube defects in the unborn child (World Health Organization, 2014)

Apart from pregnancy monitoring, health care providers at ANC usually midwives, also play

a role in providing nutrition advice to pregnant women during ANC appointments. Studies on whether midwives provide nutrition education in United Kingdom (D.J Lee, 2012) and Sweden (A.L. Wennberg, 2014) have revealed that most midwives face challenges in providing dietary advice and especially to pregnant women with medical conditions such as diabetes and although they acknowledge that it is part of their duty (J. Arrish, 2016). Midwives have attributed this difficulty to lack of adequate training on Maternal Infant and young Child Nutrition (MIYCN), limited staffing and time constraints.

The Kenya Food and Nutrition Security Policy (2018) adopts a lifecycle approach which focuses on the health and nutrition needs of individuals through six stages of human development and physiological lifecycle. The nutritional needs change with different physiological states and development in terms of specific nutrients, types and amounts and varieties of foods. Pregnancy is the first stage of the lifecycle where promoting optimal maternal nutrition through proper nutrition advice, nutrition education and counseling results to a healthier pregnancy and healthy infants. This sets stage for better nutrition status through the rest of the lifecycle.

When pregnant women enroll for ANC early in pregnancy and make several visits in the course of pregnancy, the health care givers are able to conduct assessments and identify risks and also educate women on maternal nutrition that is necessary for better pregnancy outcomes. In Kenya all public health facilities have an antenatal care clinic for pregnant women. The Kenya maternal nutrition guidelines topics include: Energy and protein requirements in pregnancy; frequency of meals; micronutrient requirements in pregnancy in terms key micronutrients like iron, zinc, iodine, vitamin A, calcium and folic acid. These are covered in terms of food sources rich in the micronutrients, their importance, consequences of deficiency, risk factors to deficiency, dietary management and micronutrient supplementation programs available. Nutritional assessment covers for: Anthropometric assessment for weight and Middle upper arm circumference measurements on every visit; Dietary assessment to assess food patterns, frequency and variety and factors contributing to inadequate intake and biochemical profile for blood sugar, evaluation for Anemia and urinalysis to check for proteinuria (MOH, 2018)

The Kenya

guideline on focused Antenatal care has adopted the WHO recommendation of at least four antenatal care visits by all pregnant women. Despite this recommendation, a 2008-2009 survey conducted on health facilities in Kenya showed that 92% of pregnant women went for antenatal care at least once but only 47% went for four or more visits as recommended by World Health Organization. The low uptake of at least four ANC visits robs women of the opportunity

to learn about their nutritional needs.

## **1.2 Statement of the Problem**

Malnutrition in all its forms affects women all over the world but its effects are worse during pregnancy period. Malnutrition is influenced by factors like socio-demography, nutritional knowledge and dietary practices of pregnant women (Beyene, 2013)

In an antenatal set up, provision of nutrition education is crucial in improving the knowledge of pregnant women about maternal nutrition

Even though many studies have been conducted in the area of maternal health, no study has been conducted to assess nutritional knowledge and dietary practices of pregnant women receiving nutrition education while attending antenatal care in the study site. The lack of comprehensive information posed the need for this study given that Mbagathi hospital is a level five facility that serves an urban catchment area with a population of 3.1 Million people (KDHS, 2014)

## **1.3 Justification of the study**

Ante natal is important in ensuring the success of a pregnancy. Nutrition education is critical in promoting good nutrition status during pregnancy and it influences maternal and child health outcomes. With adequate knowledge on nutrition during pregnancy, pregnant women can improve their dietary intake to provide proper nourishment to themselves and the developing baby. This study will be useful for the Government of Kenya in implementing policies that will improve provision of nutrition education to pregnant women. The findings of this study will also suffice as a baseline for improving the quality of the National guidelines on maternal nutrition.

## **1.4 Aim of the study**

The study aimed to contribute towards improvement of provision of nutritional education at ANC clinics in Kenya. Using the results of this study, the existing policies, guidelines and protocols of nutrition education in pregnancy can be improved to enhance the quality of maternal nutrition care.

## **1.5 Purpose of the study**

The purpose of the study was to generate data on the nutritional knowledge and dietary practices of pregnant women receiving nutrition education while attending antenatal care at Mbagathi

hospital Nairobi county The generated knowledge shall help the health managers and other stakeholders to strengthen nutrition education component at ANCs so that pregnant women can gain sufficient knowledge on how to stay healthy during pregnancy. The new knowledge can be operationalized to create awareness on barriers that hinder uptake of nutrition knowledge.

## **1.6 Objectives of the study**

### **1.6.1 Main Objective**

To determine the nutritional knowledge and dietary practices of pregnant women receiving nutrition education while attending antenatal care at Mbagathi hospital Nairobi County

### **1.6.2 Specific objectives**

1. To describe the socio-demographic characteristics of the pregnant women receiving nutrition education while attending antenatal care at Mbagathi hospital in Nairobi County
2. To determine the level of nutritional knowledge of the pregnant women receiving nutrition education while attending antenatal care at Mbagathi hospital
3. To assess the dietary diversity score of the pregnant women receiving nutrition education while attending ANC at Mbagathi hospital
4. To assess the nutrition status of the pregnant women receiving nutrition education while attending Antenatal care at Mbagathi hospital, Nairobi County
5. To determine the relationship between Antenatal care attendances, nutritional knowledge and dietary practices of the pregnant women at Mbagathi hospital in Nairobi County

## **1.7 Research Questions**

1. What socio-demographic characteristics influence the nutritional knowledge, dietary diversity and nutrition status of pregnant women receiving antenatal care at Mbagathi hospital?
2. What is the association between the level of nutrition knowledge and dietary diversity of pregnant women in Mbagathi hospital?
3. Is there a relationship between dietary diversity score and the nutrition status of pregnant women attending receiving nutrition education at Mbagathi hospital?
4. Is there a relationship between ANC attendance, nutritional knowledge and dietary practices of the study population?

## **CHAPTER TWO: LITERATURE REVIEW**

### **2.0 Introduction**

This chapter will focus on overview of ANC, nutrition education in pregnancy, nutrition knowledge of pregnant women, diet quality of pregnant women, role of nutrition in pregnancy and nutritional requirements in pregnancy. It also presents the conceptual framework and gaps in knowledge of the reviewed literature.

### **2.1 Overview of Antenatal Care**

To assure the best possible health of the mother and her fetus, various aspects such as screening, counseling, education and treatment are important. This systematic and careful assessment is called antenatal care. In the 20th century, prenatal program was first established in Europe and was mainly for poor women living in difficult conditions and its objective was to improve maternal and prenatal outcomes (Sikorski J, 1996). Over time, aspects of screening for medical problems were introduced to detect medical problems during pregnancy. As the health status of the mothers and pregnancy outcomes improved, other developed countries adopted antenatal care as much as there was no evidence of its exact benefits (Gitonga, 2017).

Over the past two decades, there has been much progress made to increase access to and use of ANC. Quality of care has also improved leading to a reduction in deaths as a result of pregnancy. Deaths because of complications that come about because of pregnancy are still very high. In 2015, about 303,000 women died because of pregnancy related complications, over 2.5 million babies died between 0-28 days after birth and 2.6 million babies were born dead. An estimate of about 64% of expectant mothers received prenatal care at least four or more in the course of pregnancy (World Health Organization, 2016).

In the most recent ANC guidelines by WHO, the number of ANC contacts have been increased

from 4 to 8 contacts throughout pregnancy. Current evidence indicates that at least eight contacts can reduce perinatal deaths by up to 8 per 1000 births compared to the 4 visit model. In the new model, WHO recommends the first visit on or before 12 weeks of pregnancy then follow up visits, taking place at 20, 26, 30, 34, 36, 38 and 40 weeks respectively (WHO, 2016). WHO recent guideline also outlines the care that pregnant women should receive during each appointment. Components include; education and counseling on healthy diet and optimal nutrition, prevention of excessive weight gain, dangerous lifestyle behavior e.g. use of illegal drugs and smoking; treatment and prevention of Malaria and HIV/AIDS; vaccination for tetanus and



blood screening; fetal ultrasound; and management of common pregnancy symptoms such as nausea, back pain and constipation. The most recent recommendation concerning nutrition in pregnancy is the daily IFAS to prevent iron deficiency anemia, puerperal sepsis, low birth weight, and preterm birth.

Antenatal care is an important tool for monitoring pregnancy and this can enable early detection and prevention of risks in the course of pregnancy, many women in developing countries do not use this service. According to KDHS (2014) about 6 in 10 women achieved at least four or more pre-natal visits, though, about only 20% had their first visit within 12 weeks. In a study to determine utilization of ANC in Delhi, 20% of the study population did not avail to any ANC services at a health facility (Ghosh-Jerath, Devasenapathy, Singh, Shankar, & Zodpey, 2015).

In a similar study to determine factors affecting ANC service utilization in Hadiya Zone Ethiopia, it was found that 86.3% of the study population used antenatal care. As much as over 80% used these services, six in ten women sought ANC in the second trimester. 41.4% of the women attended less than four visits when the recommended number is a minimum of four. (Abose, Woldie, & Ololo, 2010). ANC service utilization in Africa is generally poor with studies showing that most women seek ANC in the second semester and thus are usually unable to attain a minimum of four visits as recommended by WHO. These indicate missed opportunities for interventions like IFAS supplementation which is vital in pregnancy.

Components of nutrition education in pregnancy in the Kenya national antenatal guidelines include; education about healthy eating and prevention of excessive weight gain by keeping physically fit. Excessive weight gain can lead to a more than normal birth weight and other complications, in undernourished pregnant women, the focus of nutrition education is on improving daily intake of body building and energy giving food to minimize the risk of small for gestational babies. Nutrition education also emphasizes on the importance of multiple micronutrient supplementation. Some of the micronutrient supplements given to pregnant women during ANC visits include daily oral Iron and Folic acid which help to prevent against Iron deficiency Anemia, preterm birth, low birth weight and puerperal (www.who.int) Expectant mothers with a poor dietary calcium intake are supplemented with calcium to minimize risk of pre-eclampsia, and maternal bone demineralization and Vitamin A supplementation in populations where vitamin A deficiency is a severe to prevent night blindness (WHO, 2016) Pregnant mothers addicted to caffeine are usually advised to reduce caffeine intake to minimize pregnancy outcomes such as stillbirths and low birth weight. (WHO, 2016)

In June 2013, the government of Kenya introduced free maternal health services that

included antenatal and delivery care in government health facilities. The aim of this initiative was to reduce neonatal and maternal deaths and also increase antenatal care attendance (Malachi Arunda, 2017)

## **2.2 Nutritional knowledge of pregnant women**

When pregnant women are knowledgeable about nutrition during pregnancy they are likely to make appropriate food choices and also reject food taboos and believe about food that have not been scientifically proven.

According to Sz wajcer et al (2005) Pregnant women acknowledged that diet is important for growth and development of the fetus and are more likely to be concerned about their nutrition by seeking nutrition advice and modifying their diets. As much as there is a significant association between nutritional knowledge and maternal dietary behavior and use of supplements, sometimes pregnant women do not use their knowledge to improve nutrition dietary practices.

In a study to explore the level of nutritional knowledge of Australian pregnant women, it was found that the women exhibited high knowledge about food safety, diet health relationship and foods rich in nutrients. They exhibited low level of knowledge on topics including making everyday food choices, recommended daily intakes of five food groups, serving size, supplements during pregnancy, and importance of key nutrients in pregnancy (Bookari, Yeatman, & Williamson, 2016). In the same study, a high per cent age of the women were not aware of how much calories a gram of fat contains and also foods source high or low in fats, they also did not know the salt content of bread.

In a study in Ethiopia to assess pregnant mothers level of knowledge on Maternal Nutrition in Ethiopia, more than half of the study sample could not define the word "food". This was attributed to lack of access to information about nutrition during pregnancy since most of them were housewives. Even though they portrayed high knowledge about importance of food during pregnancy, they had limited knowledge about food source of body building, energy giving and minerals and vitamins sources. The low knowledge level of common food sources could have been due to low access to nutrition information and low socioeconomic status of the study population.

In the United States, Down set al. (2014) sought to find out how knowledgeable pregnant women were with regard to nutrition behaviors in pregnancy that could cause excessive maternal weight gain by conducting an FGD and semi-structured interviews with 30 pregnant women of the 30 participants, 65% of participants reported that many women have information on what to eat

sparingly in pregnancy, but are unaware of how much weight should be gained during pregnancy. A quarter of the participants felt they lacked adequate knowledge on healthy eating and sought information from online sources as much as some internet information lacked credibility hence was not reliable. Previously pregnant women reported feeling inadequate of nutrition information in their subsequent pregnancies because the health care providers had an assumption that they were already informed.

Shu et al. (2013) examined how knowledgeable pregnant women are on aspects of nutrition and weight gain during pregnancy. Over a half of the study population were overweight, and 75% of the overweight women were unaware of the recommended weight gain and lacked nutritional knowledge on how to prevent excessive weight gain. Only one third of women had knowledge that healthy nutrition was key to safe weight management during pregnancy. Furthermore, 51% of the women were aware that excessive weight gain in pregnancy could lead to complications such as gestational diabetes.

In Australia, Bookari et al. (2016) conducted a research on 326 expectant women in order to determine their knowledge of recommended maternal weight gain as per gestational age of pregnancy and dietary practices. At least 60% of the study population was unaware of the Australian Dietary Guidelines in pregnancy, indicating that there is unfulfilled need with regard to knowledge of maternal nutrition education. In the same study, over 50% of the participants were in agreement that being aware of how much weight should be gained was important and over 60% of the study population were informed of the appropriate weight to be gained; however, only 27.6% could accurately state their expected weight gain according to their body mass index. Bookari et al.'s (2016) findings agree with Downset al. (2014) that many women have information about what not to eat during pregnancy, but are not aware of recommended weight gain.

### **2.3 Dietary intake of pregnant women**

Good nutrition during pregnancy promotes maternal health and fetal growth. According to Rhodri et al (2010), birth outcomes such as preterm delivery, infants born with birth defects and health problems e.g. pre-eclampsia have been greatly linked to maternal diet before conception and after conception (Olafsdottir et al, 2006) and gestational diabetes (Thomas et al 2006).

Nutrient intake in pregnancy plays a role in cognitive development and allergic disorders in childhood (Nurmatov et al, 2011).

A study conducted in Spain to determine the dietary intake of pregnant women, it was found that 74% of the study population excessively consumed total fats even though the main

source however, was Mono Unsaturated Fatty Acids. Nevertheless, Saturated Fatty Acid intake was also noted to be high. In terms of sociodemographic, Latin American, young women with less education had the highest proportion of inadequate unsaturated fatty acid and folate after taking supplements because of low intake. With regard to intake of energy giving foods from starches, a higher per cent age of the Spanish women showed inadequate intake. Almost all women in the study had poor vitamin D intake according to the reference values. When taking into account total intake (food plus supplements), the percentage of inadequacy observed was still elevated (Rodríguez-Bernal et al., 2013).

Consuming saturated fat and sugar excessively and lowering consumption of starches has been linked to adverse maternal and infant health outcomes. Expectant women should thus be advised on reducing fat intake, especially saturated and increase intake of unsaturated fatty acids as well as increase intake of carbohydrates from starchy foods such as legumes

Lundqvist et al (2014) compared the dietary intake of pregnant women in early gestation and non-pregnant women found pregnant women reported less intake of total energy, meaning that pregnant women did not adhere to the recommendation on increasing energy intake during early pregnancy. This was however speculated to be due to nausea commonly experienced during the first trimester.

With regards to micronutrients, the dietary intake of vitamin D, iron and Folate among the early pregnant mothers was low. Folate plays a significant role in the fetal synthesis of DNA and RNA, low levels of Folate places risks for congenital malformations in the fetus and the other pregnancy complications.

These findings by Lundqvist et al (2014) on early pregnant women dietary intake contrasts on macronutrient intake on a study done to assess the dietary intake of pregnant mothers in Sheffield, United Kingdom. The participants were pregnant mothers in their early trimester of pregnancy. In this study, intakes of protein, energy and total fat were relatively higher. Intake of specific nutrients like vitamin C, vitamin B2, Folate and calcium was lower in the Sheffield women compared to previous studies in the region. Over 60% of the Sheffield women did not to reach the EAR (Estimated Average Requirement) for energy, Folate, vitamin B2, and iron while 40% failed to meet the EAR for calcium (Mouratidou, Ford, Prountzou, & Fraser, 2006).

A study on Dietary intakes and dietary patterns among expectant women in Northwest China, it was found that the participants' intake of fats was high; just like in other previous studies was consistent with previous studies in China (Liu FL et al, 2015). The researchers attributed this to change in lifestyle with preference for animal source foods. Blumfeld et al, 2012 also

noted an increase in fat intake above the recommendations among pregnant women in the western countries. Excessive consumption of fats leads to too much weight gain which further influence maternal and child's health, predisposing them to the risk of cardiovascular diseases, diabetes, obesity, and metabolic syndrome (Liu FL et al, 2015).

In the same study, vitamin A dietary intake was noted to be lower than the recommended intake among 70% of the study participants. This indicates that majority of pregnant women in the study area had limited access to vitamin A containing foods. A previously conducted study in China, 70% of total vitamin A was derived from plant sources, however, in the study by Liu et al, 2015 meats and vegetables accounted for 67.1% and 19.6%, respectively. This suggests a gradual change in dietary habits overtime among the study population. Dietary intake of Folate was also low and this was supported by biochemical results of low Folate concentration among women in the study area (Dang et al, 2014). Among pregnant women in developed countries, it was found that Folate intake was below the recommended (Blumfield 2013), indicating that poor dietary folate intake is a universal problem.

Still in the same study by Liu et al (2015) dietary intake of Calcium was found to be relatively higher although still below the recommended intake of any gestational period. Zinc intake was also below the recommended intake. It was speculated that absorption of calcium and zinc could be impaired by phytates that are present in the frequently consumed foods such as cereals. Iron intake was significantly higher than average and closer to recommended intake for the third trimester. These results differ from the results obtained from other developed countries where prevalence of iron deficiency is high in pregnancy.

Generally, dietary intake of pregnant women is below the recommended intake. Counseling on sufficient energy intake during the early phase of pregnancy should be emphasized. Pregnant women also need to be counselled on excessive energy intake to prevent excessive weight gain. Low intake of micronutrient has also been noted with special regard to Folate, vitamin D and iron. Nutrition educations on the importance of micronutrients need to be emphasized. Pregnant women should be encouraged to start ANC early in pregnancy in order to benefit from Folate supplementations.

Dietary diversity should be practiced by pregnant women in order to reap maximum benefits from locally available foods in order to meet the extra nutrient requirements of pregnancy. The literature reviewed above is limiting in the dietary intake of body building foods. Further studies need to be done on consumption of proteins in pregnancy.

## **2.4 Research Design and methodology**

Research study design is a framework, or a set of methods and procedures that are used for collection and analysis of data on variables specified in a particular research problem. The type of study design a researcher uses to answer a particular research question is determined by the goal of research, the nature of question and the availability of resources. The design of a study can affect the validity of its results, so it is important to understand the different types of study designs and their strengths and limitations.(Aggarwal, 2018).

Study designs are classified into descriptive(non-analytical) and analytical study designs. Descriptive studies merely describe data on one or more characteristics of a group or individual. They do not answer questions or establish causal relationships. Examples of descriptive studies include case series, case reports and cross sectional surveys (cross sectional surveys can also be analytical). A descriptive study can be a survey of dietary habits of pregnant women(Aggarwal, 2018).

Analytical studies are studies where the investigator attempts to test a hypothesis or establish a causal relationship between variables. In analytical studies, the investigator assesses the effect of an intervention or exposure of an outcome. This study is classified into observational and experimental studies. Observational studies are where the investigator documents a naturally occurring relationship between the exposure and the outcome. The researcher does not do any intervention and the exposure has already been decided naturally e.g. comparing the antenatal dietary habits of mother with normal and low birth weight babies. The exposure must precede the outcome(Aggarwal, 2018).

Observational studies can either be descriptive or analytical. Examples of observational studies include cross sectional studies, case control studies and cohort studies

A cohort study is where a study takes a forward direction, the investigator starts with determining the exposure to a risk factor and then assesses whether the outcome occurs at a future point in time eg a researcher can follow a group of smokers and a group of non-smokers to determine the incidence of lung cancer in each(Walwyn, 2016).

In cases control studies the researcher takes a backward direction, the investigator starts by determining whether the outcome is present and then traces the presence of prior exposure to a risk factor eg an investigator can identify a group of normal birthweight babies and a group of low birth weight babies and then ask the mothers about their dietary habits during pregnancy.

In interventional studies, the investigator actively performs an intervention either in all or

some participants in a study eg administration of a drug or a vaccine or introduction of an educational tool(Walwyn, 2016).

The most commonly used study design to assess maternal nutrition in pregnant women is cross sectional study design. A study in Ghana to assess nutritional knowledge and dietary habits among pregnant adolescents attending antenatal clinic used a cross sectional study design(Prince Kubi, 2021).


A cross sectional descriptive survey was used to collect data on knowledge and attitude of pregnant mothers towards maternal dietary practices during pregnancy in Yaounde(al, 2016)

In a study to assess health and nutrition knowledge attitude and practices of pregnant women attending and not attending antenatal care in Western Kenya, a cross sectional survey was used(Nandita Perumal\*, 2013).

This study used a descriptive cross sectional study design to describe the characteristics of pregnant women attending antenatal clinic at Mbagathi hospital

Unlike case control studies and cohort studies where study participants are selected based on outcome or exposure respectively, in cross sectional studies, participants are selected based on inclusion and exclusion criteria and data is collected at a specific period in time. This study does not involve manipulating variables and allow the researcher to look at numerous characteristics at once and also provide information about what happens in a current population

Descriptive cross sectional studies are inexpensive and fast, the researcher can collect data on different variables at once, they can also provide a useful springboard for further research. Nonetheless, this study design does not establish causal relationships, so it does not allow for conclusion about causation (Levin, 2006).



**Fig 1: Illustration of research study design**

## 2.4 Gaps in knowledge

There is a gap in knowledge about why pregnant women are unable to translate nutritional knowledge into dietary practices.

This data implies a gap in knowledge on the effectiveness of the nutrition education gained in the process of ante natal care or there are other factors that prevent women from fully utilizing the knowledge gained at the ANC to improve dietary practices.


## 2.5 Theoretical Framework

This study will adopt the Knowledge-Attitude-Behavior Model. Knowledge is instrumental when a health related behavior change is required. The model proposes that as one gains more knowledge in a health behavior domain, change in attitude is initiated. Over the same period of time changes in attitude accumulate resulting in behavior change. The most essential resource in this model is accumulation of knowledge which is done by provision of information; however, this model does not specify the process by which change of behavior occurs.

Nutrition education is a widely used strategy to improve nutrition status of expectant women seeking prenatal care in government run hospitals especially in developing countries. When pregnant women are knowledgeable about nutrition during pregnancy they are likely to make appropriate food choices and also reject food taboos and believe about food that have not




been scientifically proven. When women attend several ANC visits, they exhibit better nutritional knowledge, attitude, and practices compared to those not attending more so in developing (Girard & Olude, 2012). Several studies reveal that pregnant women are more receptive to nutrition information. Pregnant women need the "how-to" knowledge since they are already motivated. This kind of knowledge is instrumental to help them adopt a healthful diet that will have positive consequences in the outcome of the pregnancy.



**Figure 1: Knowledge-Attitude-Behavior (KAB) model**

## 2.6 Conceptual Framework of the study variables



**Figure 2: Conceptual framework derived from study objectives**

## **CHAPTER THREE: RESEARCH DESIGN AND METHODOLOGY**

### **3.1 Study Design**


A descriptive cross-sectional study design was used in this study. In a cross-sectional study design, the researcher measures the outcome and the exposure in the study participants at the same time and may also study their associations. The participants are selected based on the inclusion and exclusion criteria set for the study. Data was collected in the month of October and November 2021. The study sought to find out the Nutritional knowledge and dietary practices of pregnant women exposed to nutrition education while receiving antenatal care. A structured interview questionnaire was used for data collection.

### **3.2 Study site and Population**

#### **3.2.1 Study site**

This study was done at Mbagathi hospital in Nairobi County specifically at the Antenatal Care Clinic. Mbagathi Sub-county hospital is a public level 5 health facility under the County government of Nairobi Department of Health service. The hospital serves an urban catchment area with a population of 3.1 Million of which 22% live below the poverty line (KDHS, 2014).

Services offered at Mbagathi include; Medical training, special clinics, family planning, immunization, antiretroviral therapy, HIV counseling and testing, curative in-patient and outpatient services among others.



**Figure 3: Map of Kenya showing Mbagathi Hospital**

### 3.2.2 Study population

The study population consisted of pregnant women receiving nutrition education while attending antenatal at Mbagathi Hospital between the periods of 1st October 2021 to 31st November 2021.

### 3.3 Sample size

#### 3.3.1 Sample size determination

The sample size was calculated using the Fischer et al., (1991) formula as follows:

$$N = (Z^2pq) / d^2$$

Where:

N- Desired sample size

z-The standard normal deviation set at 1.96 of the 95% confidence interval

P-Prevalence of antenatal care attendance at Mbagathi hospital is at 85% (MOH, 2015)

q- Proportion of ANC attendance is  $(1-p) = (1-0.85) = 0.15$

d-The degree of accuracy desired set at 0.05 significance


Therefore,  $N = (1.96)^2(0.85 \times 0.15) / 0.052 = 195$

The study sample was therefore 195 pregnant women

### 3.3.2 Sampling procedure

Mbagathi Hospital was purposively selected since it is the largest hospital in the sub County that serves the rest of the population in the County thus more representative.

The sample for the study was selected using systematic random sampling



**Figure 4: Sampling frame for study site and study population at Mbagathi hospital**

### 3.3.3 Inclusion criteria

Participants included in this study were pregnant women receiving nutrition education while attending antenatal care visit at Mbagathi hospital who voluntarily consented and agreed to participate in the study

### **3.3.4 Exclusion criteria**

Pregnant women with ill health, complications, abnormal fetus, and those who did not voluntarily consent to the study

### **3.4 Data collection tools and procedure**

A semi-structured questionnaire was used for data collection

#### **3.4.1 Study Questionnaire**

The researcher developed a semi structured questionnaire to be used for data collection. It was divided into three parts: The first part introduced the questionnaire and briefly explain its purpose. The second part gave instructions on how to answer questions and part three covered questions regarding the independent variables: socio-demographic context, test of knowledge about nutrition in pregnancy and dietary practices of the pregnant women. Coded options were given for the questions.

##### **3.4.1.1 Socio-demographic characteristics**

Questions in this section sought to determine the socio-demographic context of the pregnant women. These included; age, marital status, education status, average monthly household income occupation, gestational age, religion and family size. This information enabled the researcher to understand the background of the respondents and also to find associations with other study variables.

##### **3.4.1.2 Test of nutritional knowledge**

Maternal nutritional knowledge was assessed using a set of questions that comprised 15 multiple choice questions that captured their understanding of various aspects of nutrition during pregnancy. This questions reflected the topics covered during nutrition education sessions women as stipulated in the Kenya national guidelines for antenatal care during .The test assessed the mother's knowledge on: variety of foods; amount of food during pregnancy; energy sources; micronutrient supplement for pregnant women, necessity for supplementation and duration; micronutrients, iron and calcium and consequences of deficiency, fetal complications of maternal under nutrition; knowledge on food safety and lifestyle behaviors to avoid during pregnancy.

The respondents selected a response to each question among the choices given. The questions were marked on a scale of 1-15. Each correctly answered question was awarded one mark. All the scores for the right answers were summed and a total score out of fifteen given. Respondents who correctly answered at least ten and above were scored as having adequate nutritional knowledge; those who correctly answered between nine and six questions were scored as having average knowledge and those who scored five and less had poor knowledge (MOH, 2018)

### **3.4.1.3 Dietary practices**

The test will assess the respondents' dietary adequacy, frequency of consumption and dietary diversity. 24-hour recall Dietary practices was assessed using a Dietary Diversity Questionnaire for Women of Reproductive Age (DDS-W) and a questionnaire

Dietary Diversity as an indicator of the quality of the diet and reflected the consumption of individual women. It sufficed as a proxy for macro and micro nutrient dietary adequacy. The study used the FAO recommended cut-off points for Minimum Dietary Diversity for Women (MDD-W) which is made of ten food groups (FAO, 2016).

The assessment involved a simple count of food groups that a participant had consumed over the preceding 24 hours before the survey. The indicator of Dietary Diversity that is the Dietary Diversity Score (DDS) was calculated by summing the number of food groups consumed by the respondent over the 24-hour recall period. These food groups include: Grains (maize, wheat, millet and all products made from these and their flours); white roots and tuber (cassava, yam, sweet potatoes, beet, Irish potatoes and others not in the list); pulses (beans, peas and lentils); nuts and seeds (pumpkin seeds, groundnuts, sunflower seeds, chia seeds, sesame seeds and others); Dairy (fresh milk, fermented milk, yoghurt, cheese); Meat, poultry and fish; Eggs; Dark green leafy vegetables (cassava leaves, pumpkinleaves, amaranth, managu, kales, spinach); other vitamin A rich fruits and vegetables (ripe mangoes and papayas, oranges, passion fruits, melons carrots, pumpkins other vegetables (fresh peas, green beans, tomatoes, cucumber, zucchini, onions, mushrooms) and other fruits (wild fruits, ripe bananas, guava, figs, lemon, lime)

Participants who had consumed at least five of the ten food groups in the 24-hour recall were considered to have met the MDD-W

The 24-hour recall questionnaire was also used to assess the actual intake of individual foods. The respondents were asked to recall and report all foods and beverages consumed over the preceding 24 hours. The recall started with the first thing the respondents ate in the

morning until the last food consumed the next morning. The steps involved in the 24-hour recall involved asking the respondents to give a list of all foods and beverages consumed during the previous 24 hours, they were then probed for foods probably forgotten during the listing, the time each food was consumed, the quantity consumed and anything else that was consumed.

### **3.5 Recruitment and Training of Research Assistants**

The study recruited five assistants with at least a certificate in Nutrition and Dietetics and experience in data collection. They administered the questionnaires and carried out direct observation. Prior to the survey, the researcher trained the assistants. The training sessions covered the purpose and objectives of the study; a thorough review of each item on the questionnaire; how to conduct an interview (role play was used for the enumerator to practice questionnaire administration) and the expected code of conduct of a researcher during an interview. After the training session, the researcher and the assistants pre-tested the questionnaire, modifying it accordingly and thereafter prepared a work plan for the execution of the study.

### **3.6 Pre -Test of the Data Collection Tools**

The questionnaire was pre-tested at Westland Health Center. The purpose of pre-testing was to check if the questions was clear, could be understood by the respondents and if they aligned with the purpose and objectives of the study. Ten percent of the study sample formed the sub sample for pre-testing the questionnaire. The observation checklist was tested on the care providers at the ANC clinic. The researcher explained the purpose and objectives of the study to the participants of the pre-test activity and obtained their consent to participate in the study. The questions were read out to the respondents. After the end of the session, the research team held a meeting to share experiences and any challenges encountered and how they could be resolved in order to make the actual study run smoothly. Screening of the filled in questionnaires was done to assess for completeness. The research went through the recommendations made by the pre-test sample and used them to improve the questionnaire.

### **3.7 Ethical and Human Rights Consideration**

Before commencement of the study, the researcher obtained approval of the study from University of Nairobi and Kenyatta National Hospital Ethics and Research Committee (UoN-KNH ERC). The researcher obtained permission and clearance to carry out the study from



the In-Charge at Mbagathi hospital. Each respondent received an explanation about the study and informed consent obtained before commencing interviews. The consent form included information such as; the title, purpose and objectives of the study, invitation to participate in the study, benefits and risks of participating and an assurance of confidentiality in handling all the information throughout the study. The questionnaire was coded to ensure anonymity of the respondents.

### **3.8 Data analysis and Quality Assurance Procedures**

All data collection tools were pretested to ensure clarity and alignment to the study objectives. The principal investigator trained the research team on all study procedures, how to administer the study tools, interviewing techniques and how to collect good quality data. At the end of each interview session, the questionnaire was screened to ensure they were completely filled by the respondent. Before data analysis, all data was cleaned and in the event of any missing data, the respondents were contacted for clarity.

Data entry was done using SPSS version 20 software. The socio-demographic characteristics were described using statistics such as mean mode, frequencies and standard deviations.

Inferential statistics: Fischer's Exact, Confidence Interval, P-Value, Spearman's Correlations were employed for data analysis.

## CHAPTER: FOUR RESULTS

### 4.1 Socio-demographic characteristics of the respondents

About 69% of the study participants were within the age category of 21-30 years, and 23% were over 30 years while 8% of the participants were below 20 years. The mean age was 27 ( $\pm 5.38$ ) years with a range of 17 to 41 years.

Majority of the respondents (79%) were married while 21% were single or not married. Nearly half (43%) of the participants had attained secondary education, 30%, 17% had completed only primary education, and 9% had obtained college and university education. Majority of the study participants (35%) were housewives, 31% made a living through self-employment, 12% were salaried workers while 5% were casual laborers. Slightly below half of the respondents (46%) had not given birth before (first pregnancy), 32% had one child, and 10% of the respondents had 3 to 4 children.

Majority of the respondents (69%) had up to 3 persons living within a household, 24% of respondents had a household of between 4 to 5 persons while 8% had a household size of over 5 persons. The mean household size was 3.18 ( $\pm 1.47$ ).

About a quarter (25%) of the respondents earned an income of between Kshs 5000-10000 while 18% and 15% of the study group earned an income of between Kshs 25001-50000 and Kshs 10001-25000. Only 6% of the women earned less than Kshs 5000 while 22% did not know how much income they earned.

About 32% spent between Kshs 5,000-10,000. Only 3% of the household spent over Kshs 50,000 a month, 24% of the respondents were unable to state how much money they used on expenses on a monthly basis, another 2% refused to disclose their monthly expenditure. Concerning religion, the most predominant religion (65%) among the respondents was protestant while 34% were catholic and 1% were Muslim and traditionists.

**Table 1: Demographic characteristics of the respondents attending antenatal care at Mbagathi Hospital**

Demographic characteristic	n=195	Number	Percent
<b>Age</b>			
21-30 years		134	69
below 20 years		16	8
over 30 years		44	23
<b>Marital status</b>			
Married		155	79

<b>Demographic characteristic</b>	<b>n=195</b>	<b>Number</b>	<b>Percent</b>
single/never married		40	21
<b>Education status</b>			
Secondary education		84	43
College		59	30
Primary education		34	17
University		17	9
No education		1	1
<b>Occupation</b>			
housewife		69	35
self employed		61	31
Salaried employee		30	15
other(specify)		26	13
Casual laborer		9	5
<b>Number of children</b>			
0		90	46
1		63	32
2		23	12
3 to 4		19	10
<b>Household size</b>			
Up to 3		134	69
4 to 5		46	24
over 5		15	8
<b>Income level</b>			
<=5000 Kshs		12	6
Kshs5,001-10,000		49	25
Kshs 25,001-50000		36	18
Kshs 10,001-25,000		30	15
Refused to disclose		8	4
Don't know	22		
<b>Household expenses</b>			
<=5000 Kshs		26	13
Kshs5,001-10,000		63	32
Kshs 10,001-25,000		36	18
Kshs 25,001-50000		14	7
>50,000 Kshs		5	3
Don't know		47	24
Refused to disclose		4	2
<b>Religion</b>			
Protestant		126	65
Catholic		66	34
Muslim		2	1
Traditionist		1	1

#### 4.2 Gestational age and ANC attendance of pregnant women at Mbagathi hospital

Over 50% of the respondents were in their third trimester of pregnancy, 34% were in second

trimester while 7% were in first trimester. On the number of visits made, 39% of the respondents had made between 3 and 4 visits to the ANC clinic, 34% had made over four ANC visits while 29% had made 2 visits. The mean ANC visits were 3.7 ( $\pm 1.65$ ) while the minimum and maximum number of visits were 1 and 9, respectively.

**Table 2: Gestational age and ANC attendance of pregnant women attending ANC at Mbagathi hospital**

<b>Characteristic</b>	<b>n=195</b>	<b>Percent</b>
<b>Gestational age</b>		
7 to 9 months (3rd trimester)		59
4 to 6 months (2nd trimester)		34
1 to 3 months (1st trimester)		7
<b>Antenatal attendance</b>		
Up to 2		28
3 to 4		39
Over 4		33
<b>Gestational age of first ANC visit</b>		
Between 1 and 2 months		29
Between 3 and 4 months		47
At 4 months and above		24

#### **4.3 Medical profile for the current pregnancy of pregnant women at Mbagathi hospital**

About 4% of the respondents had a diagnosis of high blood pressure, 2% and 1% had a diagnosis of gestational diabetes and anemia respectively. At the time of study, none of the participants had been diagnosed with fibroids, pre-eclampsia or malaria.

**Table 3: Medical profile of the respondents of pregnant women**

<b>Medical Conditions Women</b>	<b>Yes</b>	<b>No</b>
	<b>%</b>	<b>%</b>
Hypertension	4	96
Gestational	2	98

Anemia	1	99
--------	---	----

#### 4.4 Nutrition education topics taught to pregnant women attending antenatal care at Mbagathi Hospital

Almost half of the respondents (48%) acknowledged to have been informed about the importance of nutrition education by the health worker on their every visit to the clinic, 25% were occasionally provided with information on the importance of nutrition education during visits. Slightly less than two thirds of the respondents (61%) reported to have been taught about the importance of appropriate weight gain during pregnancy on every visit to the clinic, 8% were given information occasionally. Similarly, 61% of the respondents were informed about the recommended weight gain during all sessions

About 63% were always informed of the reasons for increased nutrient requirements during pregnancy on every nutrition education session, 12% were occasionally taught. Majority of the respondents (69%) were always informed about nutrient rich dietary sources, 13% were occasionally informed. Over 65% of the respondents were regularly given information on the importance of micronutrient supplementation on every ante natal visit, 15% were occasionally taught.

About 57% of the respondents were regularly given guideline for healthy eating in every ANC session, 14% were occasionally informed. Half of the respondents (50%) reported that a health worker took them through the food preparation methods that preserve nutrients and minimize contamination on every nutrition education session while 14% were occasionally taught

Slightly more than half (54%) of the respondents were always educated on the dangers of drugs and substance use during pregnancy while 12% were occasionally taught.

**Table 4: Topics taught during nutrition education sessions at Mbagathi hospital**


	always		sometimes	not at all
	n=195	%	%	%
Importance of nutrition education during		48	25	27

pregnancy			
Relevance of appropriate weight gain during pregnancy	61	11	28
Recommended weight gain during pregnancy	61	8	31
Reasons for increased nutrient requirements during requirements	63	12	25
Locally available nutrient dense foods	69	13	18
Importance of micronutrients supplementation during pregnancy	67	15	17
Issuance of information education material guidelines on healthy eating	57	14	29
Do health care providers address food safety, hygiene and appropriate food preparation methods	50	7	43
Addressing dangers of substance abuse during pregnancy	54	12	33
precaution	45	20	35
Physical distance	46	10	44
Is weight measurement taken during each visit	96	1	3
Is MUAC measurement taken during each visit	92	1	7

#### **4.5 Level of nutrition Knowledge of pregnant women attending antenatal care at Mbagathi hospital**

Majority of the respondents (84%) had adequate knowledge on maternal nutrition, 13% had average knowledge while 13% had poor knowledge score (Figure 6).

There was a significant association between the level of knowledge and number of antenatal care visits attended. Among the women who had attended more than four antenatal visits, 76% had adequate knowledge while 9% had poor knowledge. Among those who had attended four antenatal visits, 15% had adequate knowledge while 6% had poor knowledge while among those who had attended three antenatal visits, 9% had adequate knowledge while 4% had poor nutritional knowledge (p=0.03)



**Figure 5: Level of nutritional knowledge of the study participants at Mbagathi hospital**


Level of nutritional knowledge of respondents was significantly associated with the nutrition status of the study participants. In the underweight category, 20% had poor knowledge while 6% had adequate knowledge. In the normal category, 56% had poor knowledge while 66% had adequate knowledge while in the obese category 18% had poor knowledge while 28% had adequate knowledge ( $p=0.01$ )

**Table 5: Association between the level of nutritional knowledge and nutrition status(MUAC)**

Nutrition status MUAC	poor knowledge		Adequate knowledge		chi-value	p-value
	n	%	n	%		
Underweight	15	20	7	6	9.2	0.01
Normal	42	56	79	66		
Obese	18	24	34	28		

#### **4.6 Dietary Diversity score of pregnant women attending antenatal care at Mbagathi hospital**

Slightly less than two thirds (63%) of the study population consumed at least five or more food groups while 37% consumed less than five food groups within a period of 24 hours (figure 7).



**Figure 6: Dietary Diversity Scores of the pregnant women attending antenatal care at Mbagathi hospital**

There was a significant association between Dietary diversity score and respondents' household size (Table 6). Pregnant women from household sizes of 1 to 4 members displayed adequate dietary diversity ( $p=0.03$ ).

However, there was no significant association between dietary diversity score with other socio- demographic characteristics like education status, marital status, occupation and level of income.



**Table 6: Distribution of socio-demographic characteristics by dietary diversity score**

<i>Sociodemographic aspects</i>	<i>Inadequate % %</i>	<i>DDS</i>	<i>Adequate DDS</i>	<i>X<sup>2</sup>- Value</i>	<i>P-value</i>
Education level				6.2	0.18
College	33		29		
No education/Illiterate	1		0		
Primary education	23		14		
Secondary education	36		48		
University	7		10		
Marital status				0.03	0.84
Married	78		80		
Single/never married	22		20		
Occupation				4.9	0.3
Casual laborer	7		3		
Salaried employee	12		17		
Housewife	38		34		
Other(specify)	8		16		
Self employed	34		30		
Household size				0.91	0.033
1 - 4	73		65		
> 4	27		35		
Level of income				9.1	0.17
>Kshs 50,000	5		11		
Kshs 10,001-25,000	16		15		
Kshs 25,001-50000	18		19		
Kshs5,001-10,000	29		23		
Kshs<=5000	11		3		
Refused to disclose	1		6		
Don't Know	19		24		

There was a significant association between the level of nutritional knowledge and consumption of 10 recommended food groups for pregnant women. Nutritional knowledge was significantly associated with consumption of dark green leafy vegetables (p=0.045) and also consumption of other vegetables e.g. fresh peas, cabbages, green beans, and mushrooms (p=0.008).

There was also a significant association between the level of nutritional knowledge and the mean dietary diversity score (p=0.039).


**Table 7: Association between nutritional knowledge and consumption of the recommended ten food groups and the Mean Dietary Diversity score of pregnant women attending antenatal care at Mbagathi hospital**

	Low knowledge	High knowledge	N=152	Chi-square	P-value
	%	%			
Starchy staples				2.6	0.114
Never ate	3	16			
Ate	97	84			
pulses				2	0.168
Never ate	42	57			
Ate	58	43			
Nuts and seeds				0.1	0.961
Never ate	84	82			
Ate	16	18			
dairy				0.1	0.905
Never ate	29	30			
Ate	71	70			
Meat poultry fish				0.7	0.427
Never ate	65	55			
Ate	35	45			
eggs				0.1	0.933
Never ate	77	79			
Ate	23	21			
Dark green leafy vegetables				4.1	0.045
Never ate	19	40			
Ate	81	60			
Other Vitamin A rich fruits and vegetables				0.5	0.526
Never ate	26	34			
Ate	74	66			
Other vegetables				7.2	0.008
Never ate	10	36			
Ate	90	64			
Other fruits and vegetables				3	0.085
Never ate	35	54			
Ate	65	46			
MDDS score				4.3	0.039
Never ate	19	41			
Ate	81	59			

---

#### 4.7 Nutritional status of the pregnant women attending antenatal care at Mbagathi hospital

Figure 7 represents the nutritional status of the pregnant women based on their mid-upper arm circumference measurements (MUAC). About 62% of the pregnant women had normal nutritional status with a MUAC measurement of between 22.8 - 30.57. About 27% of the women were obese with a MUAC readings of >30 cm while 11% of the women were underweight with MUAC reading of <22 cm.



**Figure 7: Nutritional status of pregnant women attending antenatal care at Mbagathi hospital based on MUAC cut-off points**


#### Key

22.8-30.57	Normal
>30.57	Obesity
<22.8	Underweight

Figure 8 shows the association between knowledge levels and nutritional status. Among the women who were underweight with a MUAC reading of > 22cm, 26% had low knowledge on maternal nutrition while 9% had high knowledge. In the normal nutritional status category, 65% had high maternal nutrition knowledge, while 48% had low knowledge. There was no difference in knowledge levels in the obese group

There was a statically significant difference between the nutritional knowledge and the

nutritional status of the pregnant women ( $p=0.018$ ).



**Figure 8: Association between the level of nutritional knowledge and nutritional status of the women attending antenatal care at Mbagathi hospital**

## **CHAPTER FIVE: DISCUSSION**

### **5.1 Introduction**

Antenatal care is a pregnancy related service provided to expectant women by health care workers to prevent neonatal deaths and maintaining the health of a woman during pregnancy. It is becoming increasingly evident that pregnancy period is an influential and teachable moment to promote nutrition among pregnant women (Arrish, Yeatman, & Williamson, 2014). Due to regular contact with health care providers, pregnant women are interested and actively seek for more information regarding maternal nutrition (Obago, 2013).

### **5.2 Socio-demographic context of the study participants**

Socio-demographic context of a study population has been found to have significant associations with variables under investigation in a study.

There was no significant association between nutritional knowledge levels and a socio-demography aspect like education status, occupation, marital status and household income. In a study to assess dietary patterns of pregnant women in Nepal also found no association between women literacy levels and knowledge of nutrition (Shah et al., 2017). However, Studies by (Ogunba&Abiodun, 2017) and (Leong Wong, ZheXin, & Lean Keng, 2018) revealed a significant association between nutritional knowledge and socio-demographic characteristics like education status, occupation and household income. These two studies indicate socio- demographic factors like high education status, play a role in that pregnant mothers with high education levels displayed high nutritional knowledge due to their ability to understand nutrition information disseminated by health workers. However, (Arrish et al., 2014) suggests that the period of pregnancy makes women more teachable and regular interaction with care (Placeholder4) (Placeholder4)providers makes them actively seek nutrition information for better pregnancy outcomes. This could explain the lack of significance between socio-demographic factors and nutritional knowledge in this study. All pregnant women irrespective of their social-demographic context are receptive of nutrition information and antenatal health care workers should put in more effort to pass information in a simple and easy to understand language coupled with practical demonstration and can also help the women understand the information better.

This study noted a significant association between the household size of the respondents with their dietary diversity score. A study to find out predictors

of adequate dietary diversity among pregnant women in Ethiopia showed that pregnant women from a family size of less than five were three times more likely to have an adequate dietary diversity than their counterparts from a family size of six and above (Kuma, Tamiru, & Belachew, 2021).

A possible explanation for the presence of significance in this study a small household has a lesser burden of food distribution compared to a large household.

### **5.3 Level of Nutritional knowledge of the study participants**

Provision of nutrition education by health care professionals is among the key components of ANC where during ANC sessions, pregnant women are equipped with knowledge on maternal nutrition which is supposed to influence their dietary practices for better pregnancy outcome. In this study, more than two thirds of the respondents had adequate nutritional knowledge

The results of this study are in agreement with a similar study in Malaysia (Leong Wong et al., 2018) where more than half of the respondents (63%) had high knowledge regarding maternal nutrition during pregnancy. About 61% of participants in a study by (Nana & Zema, 2018) scored highly on most statements to assess nutrition knowledge except for knowledge about iodine, vitamin A and iron food sources. (Mshanga, Martin, & Petrucka, 2020).

In a study by (Zerfu & Biadgilign, 2018) pregnant mothers scored poorly in nutrition knowledge questions. They had very limited knowledge on what constitutes a balanced diet, benefits of key micronutrients during pregnancy, food sources of specific nutrients and rich dietary sources of iron and vitamin A

When pregnant women are adequately equipped with knowledge regarding their nutritional needs, they are likely to improve their dietary intake and nutrition status

The level of nutrition knowledge of pregnant mothers can be influenced by many factors including their level of education of the mother, occupation, the number of ANC sessions attended, practical demonstration during nutrition education sessions, household income (Leong Wong et al., 2018), (Zerfu & Biadgilign, 2018) and (Emara, 2019).

In this study, adequate nutrition knowledge displayed by the respondents could be because of vast topics on nutrition taught during nutrition education sessions

Most of the respondents had attained secondary education, and that could have made it easier for them to receive, process and understand information disseminated to them by health workers at the clinic

#### **5.4 Dietary practices of the pregnant women attending antenatal care at Mbagathi hospital**

About 63% of the respondents had an adequate dietary diversity score because they had consumed five or more food groups of the ten food groups 24 hours preceding the survey. Over 50% of participants in studies by (Zelalem, Endeshaw, Ayenew, Shiferaw, &Yirgu, 2017a) and (Nana &Zema, 2018) had adequate dietary diversity score. However, (Aliwo, Fentie, Awoke, &Gizaw, 2019) found that only 31% had an adequate dietary diversity score. The dietary diversity of pregnant women can be affected by factors such as level of income, the level of nutrition knowledge and the household size

Education status, household food insecurity and level of nutritional knowledge were significantly associated with the dietary practice of study participants in a study by (Tsegaye, Tamiru, &Belachew, 2020). Participants who had attained secondary education were 3.4 times more likely to consume a diversified diet, women from food secure households were 1.6 times more likely to consume a diversified diet.

Delil, Zinab, Mosa, Ahmed, &Hassen, (2021) also found out that major determinants of dietary practices of pregnant women in their study were education level of both the woman and her partner, level of household income, occupation of the participant and partner and whether the participant was receiving nutrition information.

In the current study, adequate level of nutritional knowledge significantly influenced the participant's dietary practices. Pregnant women who are aware of their nutritional needs are more likely to consume more diversified diets than their counterparts with poor or low knowledge.

The mean household size of three could also have enabled for households to afford variety hence contributing to a high dietary diversity score.

#### **5.5 Distribution of nutritional knowledge and mean dietary diversity score of pregnant women attending antenatal care at Mbagathi hospital**

This study revealed a significant association between nutritional knowledge and the mean dietary diversity score for the ten food groups. For specific food groups, there was a strong association between nutritional knowledge and consumption of dark green leafy vegetables and other colored vegetables at.

In a study by Zelalem, Endeshaw, Ayenew, Shiferaw, & Yirgu, (2017b) to find out the effects of nutrition education on nutritional knowledge and dietary practices among pregnant women in Addis Ababa, the participants dietary practices improved from 46.8% to 83.7% post exposure to nutrition education. This study implies that provision of nutrition education improved the nutritional knowledge of the mothers hence an improvement in their dietary practices

(Tsegaye et al., 2020) found out that nutritional knowledge was a key factor associated with dietary practices of the respondents in the study. Mothers who had good nutritional knowledge were 1.7 times more likely to consume a diversified diet compared to their counterparts. Nutrition education provides women with knowledge on how to improve their dietary. With adequate nutritional knowledge, pregnant women are likely to make appropriate food choices and also ensure variety of nutrient dense foods in their daily food selection.

Health care providers at ANC clinics need to continuously create more awareness on the importance of adequate consumption of food different food groups. This foods groups offer a variety of important nutrients like proteins, carbohydrates, vitamins and minerals and essential fatty acids (FAO & 360, 2016) and (Perumal et al., 2013). Pregnant women need these nutrients in larger amounts more than non-pregnant women for growth and development of the baby, building immunity, brain development for the growing baby and energy provision. If pregnant mothers are made more aware of the specific functions of these nutrients, it would influence their dietary practices hence much better scores for this nutrients.

#### **5.6 Nutrition status of the study participants attending antenatal care at Mbagathi hospital**

In this study, the good nutrition status displayed by more than half of the participants could be attributed to high knowledge about maternal nutrition and adequate dietary diversity.

In a study to assess the nutritional knowledge and nutrition status of pregnant women attending antenatal care in Ethiopia, 80% of the pregnant women had a normal nutrition status and 19% were underweight (Nana & Zema, 2018)

#### **5.7 Distribution of nutritional knowledge levels by nutritional status of pregnant women attending antenatal care at Mbagathi hospital**

This study revealed a significant association between nutritional status and the level of



nutritional knowledge

When women are knowledgeable about their nutrition needs in pregnancy, they are likely to improve their nutrition practices hence better nutrition status. Some women however may have high knowledge but are either undernourished or overweight. (Beyene, 2013) Presence of a medical condition, low income level and large household size could be some of the factors hindering them from achieving optimum nutrition in (Asma, Asma-Ul-Hosna, & Ashraful, 2012 and Lundqvist et al., 2014).

### **5.8 Association between the level of nutritional knowledge and antenatal care attendance**

This study displayed a significant association between the number of antenatal visits attended with the level of nutritional (p=0.03). Study participants who had attended four or more antenatal sessions displayed high levels of nutritional knowledge. Zelalem et al., (2017b) found out that exposure to nutrition education during antenatal visits improved the nutritional knowledge of study participants from 53.9% pre exposure to 97% post exposure. These implies that nutrition education offered by health care providers during antenatal care sessions play a key role in improving nutritional knowledge of pregnant mothers. Therefore, health care providers should encourage pregnant mothers to attend at least four or more sessions. The earlier a pregnant woman initiates antenatal visits, the more knowledge the pregnant woman is likely to gain.

## **CHAPTER SIX: CONCLUSIONS AND RECOMMENDATIONS**

### **6.1 Conclusion**

In this study, majority of the respondents had adequate knowledge on maternal nutrition. The women who attended more ante natal sessions displayed high level of nutritional knowledge than the ones who attended fewer sessions. Women with high knowledge scores had adequate dietary diversity and good nutrition status. However, most of the women in the overweight and underweight category, displayed poor maternal nutritional knowledge.

### **6.2 Recommendations**

1. The Ministry of Health and partners need to continuously create more awareness and encourage women to initiate ante natal care early in pregnancy and attend follow up visits as required.
2. Particular attention has to be paid to the underweight and overweight women since they display low knowledge levels. Health care professionals should go an extra mile to find out why they are unable to make the right nutrition choices where food is concerned.
3. The Government should provide (Other than maternal nutrition guidelines used by health workers), carry home reference materials like flyers should be available for pregnant women. Food group dummies should also be used in order to improve the richness of the session.
4. Similar studies need to be carried out country wide in order to generate more data that can be used to improve the standards for maternal care in Kenya as well as to inform policy. Overall, however, pregnant women should be encouraged to start ante natal care within the first trimester of pregnancy and also ensure they attend every subsequent session to maximum benefit from the services offered.

## REFERENCE

- Abosse, Z., Woldie, M., & Ololo, S. (2010). Factors influencing antenatal care service utilization in hadiya zone. *Ethiop J Health Sci*, 20(2), 756-82. <https://doi.org/10.4314/ejhs.v20i2.69432>
- Aliwo, S., Fentie, M., Awoke, T., & Gizaw, Z. (2019). Dietary diversity practice and associated factors among pregnant women in North East Ethiopia. *BMC Research Notes*, 12(1), 166. <https://doi.org/10.1186/s13104-019-4159-6>
- Arrish, J., Yeatman, H., & Williamson, M. (2014). Midwives and nutrition education during pregnancy: A literature review. *Women and Birth*. <https://doi.org/10.1016/j.wombi.2013.02.003>
- Asma, R. S., Asma-Ul-Hosna, & Ashraful, A. (2012). Pregnancy and nutrition. *Bangladesh Journal of Medical Science*, 11(4), 267-272. <https://doi.org/10.3329/bjms.v11i4.12596>
- Bookari, K., Yeatman, H., & Williamson, M. (2016). Exploring Australian women's level of nutrition knowledge during pregnancy: A cross-sectional study. *International Journal of Women's Health*, 8, 405-419. <https://doi.org/10.2147/IJWH.S110072>
- Delil, R., Zinab, B., Mosa, H., Ahmed, R., & Hassen, H. (2021). Determinants of dietary diversity practice among pregnant women attending antenatal clinic at Wachemo University Nigist Eleni Mohammed memorial referral hospital, Southern Ethiopia. *PLoS ONE*, 16(4 April), 1610. <https://doi.org/10.1371/journal.pone.0250037>
- Emara, H. A. (2019). Effect of Nutrition Education Package on Pregnant Women Knowledge and Healthy Dietary Practice, 8(1), 386-44. <https://doi.org/10.9790/1959-0801013844>
- FAO, & 360, F. H. I. (2016). *Minimum Dietary Diversity for Women: A Guide for Measurement*. Retrieved from <http://www.fao.org/3/a-i5486e.pdf>
- Ghosh-Jerath, S., Devasenapathy, N., Singh, A., Shankar, A., & Zodpey, S. (2015). Ante natal care (ANC) utilization, dietary practices and nutritional outcomes in pregnant and recently delivered women in urban slums of Delhi, India: an exploratory cross-sectional study. *Reproductive Health*, 12(1). <https://doi.org/10.1186/s12978-015-0008-9>
- Girard, A. W., & Olude, O. (2012). Nutrition education and counselling provided during pregnancy: Effects on maternal, neonatal and child health outcomes. *Paediatric and Perinatal Epidemiology*, 26(SUPPL. 1), 191-204. <https://doi.org/10.1111/j.1365-3016.2012.01278.x>
- Gitonga, E. (2017). Determinants of Focused Antenatal Care Uptake among Women in Tharaka Nithi County, Kenya. *Advances in Public Health*, 2017, 164. <https://doi.org/10.1155/2017/3685401>
- Hijazi, H. H., Alyahya, M. S., Sindiani, A. M., Saqan, R. S., & Okour, A. M. (2018). Determinants of antenatal care attendance among women residing in highly disadvantaged communities in northern Jordan: A cross-sectional study. *Reproductive Health*, 15(1), 1618. <https://doi.org/10.1186/s12978-018-0542-3>
- Hodgins, S., & D'Agostino, A. (2014). The quality-coverage gap in antenatal care: Toward

better measurement of effective coverage. *Global Health Science and Practice*, 2(2), 1736181. <https://doi.org/10.9745/GHSP-D-13-00176>

- Kuma, M. N., Tamiru, D., & Belachew, T. (2021). Level and predictors of dietary diversity among pregnant women in rural South- - West Ethiopia: a community- - based cross- -sectional study. <https://doi.org/10.1136/bmjopen-2021-055125>
- Leong Wong, J., ZheXin, L., & Lean Keng, S. (2018). *KNOWLEDGE OF NUTRITION DURING PREGNANCY AND ASSOCIATED FACTORS AMONG ANTENATAL MOTHERS* The distribution of Legionella pneumophila Serogroup 1 genotypes in water cooling towers and water distribution systems in public buildings View project Knowledge of dengue among community in Kubang Kerian, Kelantan View project. Article in *International Journal of Public Health and Clinical Sciences*. Retrieved from <https://www.researchgate.net/publication/323941595>
- Lundqvist, A., Johansson, I., Wennberg, A. L., Hultdin, J., Högberg, U., Hamberg, K., & Sandström, H. (2014). Reported dietary intake in early pregnant compared to non-pregnant women - A cross-sectional study. *BMC Pregnancy and Childbirth*, 14(1). <https://doi.org/10.1186/s12884-014-0373-3>
- Mouratidou, T., Ford, F., Prountzou, F., & Fraser, R. (2006). Dietary assessment of a population of pregnant women in Sheffield, UK. *British Journal of Nutrition*, 96(5), 929-935. <https://doi.org/10.1017/BJN20061945>
- Mshanga, N., Martin, H., & Petrucka, P. (2020). Knowledge and Dietary Practices on Vitamin A and Iron Among Maasai Pregnant Women: A Mixed-Method Study, Ngorongoro, Tanzania. *Nutrition and Dietary Supplements, Volume 12*(September), 179-188. <https://doi.org/10.2147/nds.s264592>
- Nana, A., & Zema, T. (2018). Dietary practices and associated factors during pregnancy in northwestern Ethiopia. *BMC Pregnancy and Childbirth*, 18(1), 168. <https://doi.org/10.1186/s12884-018-1822-1>
- Nankumbi, J., Ngabirano, T. D., & Nalwadda, G. (2018). Maternal Nutrition Education Provided by Midwives: A Qualitative Study in an Antenatal Clinic, Uganda. *Journal of Nutrition and Metabolism*, 2018. <https://doi.org/10.1155/2018/3987396>
- Obago. (2013). Does the Quality of Antenatal Care Predict Health Facility Delivery Among Women in Kenya? Further Analysis of KDHS Data 2008/09. *International Journal of Child Health and Nutrition*, (February). <https://doi.org/10.6000/1929-4247.2013.02.01.3>
- Ogunba, B. O., & Abiodun, O. B. (2017). Knowledge and Attitude of Women and Its Influence on Antenatal Care Attendance in Southwestern Nigeria. *J Nutr Health Sci*, 4(2), 207. Retrieved from [www.annepublishers.com](http://www.annepublishers.com)
- Organization, world health. (2016). WHO Recommendation on Antenatal care for positive pregnancy experience. *WHO Recommendation on Antenatal Care for Positive Pregnancy Experience*, 152. [https://doi.org/ISBN 978 92 4 154991 2](https://doi.org/ISBN%20978%2092%204%20154991%202)
- Perumal, N., Cole, D. C., Ouédraogo, H. Z., Sindi, K., Loechl, C., Low, J., Oyunga, M. (2013). Health and nutrition knowledge, attitudes and practices of pregnant women

- attending and not-attending ANC clinics in Western Kenya: A cross-sectional analysis. *BMC Pregnancy and Childbirth*, 13. <https://doi.org/10.1186/1471-2393-13-146>
- Rodríguez-Bernal, C. L., Ramón, R., Quiles, J., Murcia, M., Navarrete-Muñoz, E. M., Vioque, J. Rebagliato, M. (2013). Dietary intake in pregnant women in a Spanish Mediterranean area: As good as it is supposed to be? *Public Health Nutrition*, 16(8), 137961389. <https://doi.org/10.1017/S1368980012003643>
- Rurangirwa, A. A., Mogren, I., Nyirazinyoye, L., Ntaganira, J., &Krantz, G. (2017). Determinants of poor utilization of antenatal care services among recentlydelivered women in Rwanda; a population based study. *BMC Pregnancy and Childbirth*, 17(1), 1610. <https://doi.org/10.1186/s12884-017-1328-2>
- Shah, S., Sharma, G., Shris, L., Shah, S. K., Sharma, M., &Sapkota, N. K. (2017). Knowledge on dietary patterns among pregnant women attending antenatal care checkup in Narayani hospital, Nepal. *International Journal of Community Medicine And Public Health*, 4(5), 1466. <https://doi.org/10.18203/2394-6040.ijcmph20171756>
- Tsegaye, D., Tamiru, D., &Belachew, T. (2020). <p>Factors Associated with DietaryPractice and Nutritional Status of Pregnant Women in Rural Communities of Illu Aba BorZone, Southwest Ethiopia</p>. *Nutrition and Dietary Supplements, Volume 12*, 1036112. <https://doi.org/10.2147/nds.s257610>
- Tuncalp, Ö., Rogers, L. M., Lawrie, T. A., Bucagu, M., Barreix, M., Peña-, J. P., í Asia, E. (2020). WHO recommendations on antenatal nutrition: an update on multiple micronutrient supplements, 366. <https://doi.org/10.1136/bmjgh-2020-003375>
- World Health Organisation. (2003). Working with Individuals, Families and Communities to Improve Maternal and Newborn Health. *World Health Organization*, 63. <https://doi.org/10.1017/CBO9781107415324.004>
- World Health Organization. (2014). Nutrition counselling during pregnancy. *World Health Organization*. Retrieved from[http://www.who.int/elena/titles/nutrition\\_counselling\\_pregnancy/en/](http://www.who.int/elena/titles/nutrition_counselling_pregnancy/en/)
- Zelalem, A., Endeshaw, M., Ayenew, M., Shiferaw, S., &Yirgu, R. (2017a). Effect of Nutrition Education on Pregnancy Specific Nutrition Knowledge and Healthy DietaryPractice among Pregnant Women in Addis Ababa. *Clinics in Mother and Child Health*, 14(3). <https://doi.org/10.4172/2090-7214.1000265>
- Zelalem, A., Endeshaw, M., Ayenew, M., Shiferaw, S., &Yirgu, R. (2017b). Effect of Nutrition Education on Pregnancy Specific Nutrition Knowledge and Healthy DietaryPractice amongPregnant Women in Addis Ababa. *Clinics in Mother and Child Health*, 14(3). <https://doi.org/10.4172/2090-7214.1000265>
- Zerfu, T. A., &Biadgilign, S. (2018). Pregnant mothers have limited knowledge and poor dietary diversity practices, but favorable attitude towards nutritional recommendations in rural Ethiopia: Evidence from community-based study. *BMC Nutrition*, 4(1), 169. <https://doi.org/10.1186/s40795-018-0251-x>

## APPENDIX 1: STUDY QUESTIONNAIRE

Questionnaire Number: . . . . .

Name of Interviewer: . . . . . Date: . . . . .

### SECTION A: SOCIO-DEMOGRAPHIC AND SOCIOECONOMIC CHARACTERISTICS

Please circle or fill in the most appropriate response where applicable

1. What is your age in years? \_\_\_\_\_
2. What is your marital status now?
  - A. Married
  - B. Single /Never married
  - C. Widowed
  - D. Separated
  - E. Divorced
3. What is your level of education?
  - A. No education/ Illiterate
  - B. Primary education
  - C. Secondary education
  - C. College
  - D. University
4. What is the gestational age of your pregnancy?
  - A. First trimester
  - B. Second trimester
  - C. Third trimester
5. What is your main occupation?
  - A. Salaried employee
  - B. Self-employed
  - C. Farmer
  - D. Casual laborer
  - F. Housewife
  - E. Other (specify)
6. What is the monthly income level of your household?
  - A. < Kshs 5000
  - B. Kshs 5000-10000
  - C. Kshs 10000-20000
  - D. Above Kshs 20000
7. What is your family size?
  - A. 1
  - B. 2
  - C. 3
  - D. 4 plus
8. Receiving nutrition information?
  - A. Yes (specify who gives information)
  - B. No
9. Other than antenatal clinic,

**SECTION B: TEST OF NUTRITION KNOWLEDGE**

In this section, I am going to ask you some questions about nutrition of pregnant women. Please let me know if you need me to clarify any of the questions. Feel free to ask any question you may have.

10. What do you think is the importance of adequate nutrition during pregnancy?
- A. To support the growth and development of the fetus and maintenance of the woman's own health
  - B. To support growth and development of the fetus
  - C. I don't know
  - D. Other  
(specify) í

11. How should a pregnant woman eat in comparison with a non-pregnant woman to provide good nutrition to her baby and help him/her grow?
- A. Eat a variety of foods from different food groups
  - B. Eat only what she craves
  - C. Maintain a normal serving size of food as a non-pregnant woman
  - D. Other (specify)

11. During pregnancy, it is advisable for a pregnant woman to
- A. Eat an extra meal on top of the three regular meals
  - B. Eat less food
  - C. Eat the same as before during pregnancy
  - D. Others (specify)

12. Can you name a potential problem when a pregnant woman gains too little weight?
- A. Risk of giving birth to a low weight baby
  - B. Risk of baby being born prematurely
  - C. I don't know
  - D. Other (specify) í ..

13. Name one complication that can arise when a pregnant woman gains too much weight that is above the recommended?
- A. Chances of giving birth via caesarian section increases
  - B. A



14. Most pregnant women would benefit from two a supplement type or tablet during pregnancy. Which one?

- A. Folic acid
- B. Iron and folic acid supplement
- C. Don't know

15. Can you tell me why it is important to take that supplement during pregnancy?

- A. To prevent anemia and birth defects
- B. To increase appetite
- C. Don't know

16. What is the frequency of taking the above named supplement during pregnancy?

- A. Daily throughout pregnancy
- B. Once in a week
- C. I don't know

17. How often should a baby younger than six months be breast fed or fed with breast milk?

- A. On demand, whenever the baby wants
- B. After 3 hours
- C. Don't know
- D. Other (specify) í

18. What is the importance of iron during pregnancy?

- A. Formation of blood for both the mother and baby
- B. Appetizer
- C. Energy giving
- D. Control of heartburn

19. Why should you prevent meat and other raw foods from touching other foods such as those that are cooked or ready to eat?

- A. Raw foods may contain germs which may be transferred to other foods
- B. Don't know
- C. Other

20. What should you do before washing fruits and vegetables?

- A. Wash them thoroughly before cooking or eating
- B. Peel them before washing
- C. Other
- D. Don't know

**21. Identify practices to avoid during pregnancy**

- A. Smoking
- B. Alcohol consumption
- C. Use of illegal drugs
- D. All of the above

**SECTION C: INDIVIDUAL DIETARY DIVERSITY SCORE**

Kindly let me know the foods that you consumed in the past 24 hours.(Before commencing the questionnaire, ask if the day before the interview was normal day i.e. no festivals, no feast, no fast; if YES, proceed to other questions, if NO ,continue. Include any snacks consumed

Food group	Example of locally available foods	YES=1, NO=0
1. Grains,white roots tubers and plantains	Maize wheat, sorghum,millet, breads, cassava,sweet potatoes, yams Ugali, porridge and other grains and other foods made from these	
2. Pulses	Beans, peas,lentils,chick pea, cow pea, soy bean	
3. Nuts and seeds	Sesame seeds, sunflower seeds, pumpkin seeds, groundnuts, macadamia, cashew nuts	
4. Dairy	Milk, cheese, yoghurt	
5. Meat poultry fish	Red meat, organ meats, sea food, chicken meat	
6. Eggs	From chicken, ducks, guinea fowls	
7. Dark greeny leafy vegetables	Cassava leave, pumpkin leaves, bean leaves, amaranth, manage, kales, spinach	
8. Other vitamin A rich fruits and vegetables	Ripe mangoes and papaya passion fruits, melons, carrots, pumpkin, red pepper	
9. Other vegetables	Fresh peas, green beans,tomato, cucumber okra, onions, mushrooms	
10. Other Fruits	Wild fruits, apple, ripe bananas, guava, figs, grapes, kiwi fruit, lemon, lime	

**Thank you for your time and participation in this study**

**APPENDIX 2: OBSERVATION GUIDE AT THE ANTE NATAL CLINIC  
INFLUENCE OF NUTRITION EDUCATION ON NUTRITIONAL KNOWLEDGE  
AND DIETARY PRACTICES OF PREGNANT WOMEN ATTENDING  
ANTENATAL CARE AT MBAGATHI HOSPITAL**

Name of Researcher \_\_\_\_\_

Date \_\_\_\_\_

Please observe the following practices and availability of the mentioned items at  
ANC clinic

- 1) Nutrition education topics essential in pregnancy

QUESTIONS ON KEY NUTRITION INFORMATION THAT SHOULD BE PROVIDED DURING ANC APPOINTMENTS	ANSWERS 1) YES 2) NO
1. Are pregnant women taught about the importance of adequate nutrition during pregnancy?	
2. Are pregnant women informed about relevance of appropriate weight gain during pregnancy?	
3. Are pregnant women taught about increased nutrient requirements?	
4. Do service providers mention some of the nutrient rich sources locally available suitable for pregnant women?	
5. Are pregnant women educated on the importance of micronutrient supplementation during pregnancy?	
6. Are pregnant women issued with information education materials such as leaflets, handouts or brochures on guidelines for healthy eating habits	
7. Do service providers address food safety, hygiene and appropriate food preparation methods?	
8. Are pregnant women taught about nutrition in special circumstances such as chronic diseases?	

**APPENDIX 3: GUIDELINES AND RESOURCE MATERIALS  
AVAILABLE ON MATERNAL NUTRITION**

GUIDELINES/RESOURCE MATERIALS AVAILABLE	AVAILABILITY 1)YES 2)NO	USE 1)YES 2) NO
National maternal, infant and young child nutrition policy		
Maternal, Infant and Young Child Nutrition Guidelines for Health Workers		
Take home brochure on Nutrition during Pregnancy and Lactation		

## APPENDIX 4: CONSENT FORM

### FOMU YA IDHINI YA KUSHIRIKA KATIKA UTAFITI

**KICHWA CHA UTAFITI:** UshawishiWaElimuYaLishe Bora KwenyeKiwango Cha  
MaarifaYaLishe \_\_\_\_\_ Bora \_\_\_\_\_ Na \_\_\_\_\_ Mazoea  
YaUlajiWaWanawakeWajawazitoWanaopokeaHudumaYaKlinikiYaUjaUzitoKatikaHospitali  
YaMbagathi

#### Mtafiti mkuu Na taasisi husika:

Marygorret Maloba, mwanafunzi wa shahada ya Sayansi, idara ya FSNT, chuo kikuu cha Nairobi

#### Watafiti-wenza Na taasisi husika:

1. Professor Wambui Kogi-Makau, Mkuu wa idara, FSNT, chuo kikuu cha Nairobi
2. Dr Mulwa Dasel Kaindi, Mkufunzi, Idara ya FSNT, chuo kikuu cha Nairobi

#### UTANGULIZI

Ningependakukueleza juu ya utafiti utakaofanywa na watafiti iamba owa metajwa. Kusudi la fomuhii ya idhini ni kukupa maelezo ambayo yatakuambia kuamua iwapo utashiriki katika utafiti huu au la. Kuwahuru kuulizama swali yote kuhusu kusudi la utafiti huu, lipi litakalo fanyika unapokubalika shirikikataika utafiti, madahara na faida ya utafiti, hakizakokamamshiriki wakujitolea chochote kile ambacho kina utakuhusu utafiti huu. Utakaporidhika namajibu yamaswali yako, utakuwahuru kuchagua kushiriki au la. Utakapoelewa kusudi la utafiti huu naku kubalika shiriki nitakuomba utiesahihika ikiwa fomuhii Kanunizi na zingati wakati utafiti wakia fyanika mazifwatazo:

1. Uamuzi wa kushiriki kushiriki katika utafiti wakujitolea
2. Waweza kuamua kujiondoa katika utafiti pasipokutoa sababu zote za kujiondoa
3. Kukataa kushiriki katika utafiti hakutaadhirika kizako za kupata huduma za kiafya katika hosipitali hii ama hosipitali zingine

Tutakupanaka la ya fomui ya idhini uweke kwa ajili ya kumbukumbu

Je, niendeleo? **NDIO/HAPANA**

Utafiti huu meidhinishwa na kamati ya utafiti ya Hospitali kuuya \_\_\_\_\_ Kenyatta ikishiriki ana chuo kikuu cha Nairobi. Nambari ya utafiti ni í í í í í í í í ..

#### UTAFITI HUU UNAHUSU NINI?

Watafiti waliotajwa hapombeleni wanahojiakina \_\_\_\_\_ mama \_\_\_\_\_ wajawazito. Kusudi la mahojianoni kutakajuai wapo iwapo elimu ya lishe inayotolewa katika kliniki wakati waujauzitoi

nachangiakatikakukuzakiwango cha maarifayalishenamazoeayaulajiyawanawakewajawazito. Washiriki katika utafiti huu wataulizwa maswali kuhusu mazoea ya ulaji wao katika haliyaoujauzito. Takribanwashiriki 152 walichaguliwa kushiriki kupitia njia ya bahati nasibu

**NI LIPI LITAKALOFANYIKA IWAPO UTAAMUA KUSHIRIKI KATIKA UTAFITI HUU?**

Iwapoutakubalikushirikikatikautafiti, yafwatayoyatafanyika

1. Utahojiwafaraganinamtafitiambayeamehitimuiliuwehurukujibumaswali
2. Mahojianoyatachukuakiasi cha dakika 25-35.
3. Mahojianoyatahusumaswalayalishe bora wakatiwaujauzito, nyongezayamadini ya iron na folic acid, nautumiajiwadawanapombewakatiwaujauzito

Tutakuombautopenambariyakoyasimuituwezekuwasiliananawetutakapohitajimajibukwamas waliambayoutakuwahaujajibu.Nambariyakoyasimuitatumiwatuna wale watuambaowanahusikana utafiti huu.

**KUNA MADHARA, HATARI NA USUMBUFU WOWOTE AMBAO UNaweZA TOKANA NA UTAFITI HUU?**

Utafitiwakiafyauawezakuletamaharakwamshirika.Kamawatafiti, tunatiajuhudizotekupunguzamadharayo yote.Baadhiyamadharayautafitinikupotezafaragayams hiriki.Tunakuhakikishiyakwambayaleyoteutakayoutuelezeatutayawekasiri.Tutatumianambar i fiche badalayajinalakokukutambua

Madharamenginikwambamaswalimengineyanayoulizwayanawezatibuahaliyawasiwasi.

Iwapokutakuwanamaswaliambayohautapendakujibu,unawezayaruka.Unahakiiyakukataakuho jiwaamakujibumaswalimenginewakatiwamahojiano

Kama watafiti, tunaelewakwambssivizurikuendeshamahojianopalipowatuwengi/umma, kwahivyotutafanyamahojianofaraganiambapoutakuwahurukujibumaswali.

Mwisho, mtafitiambayeatakuhojiamehitimukatikamaswalayakuendeshamahojiano

**KUNA FAIDA YOYOTE KATIKA UTAFITI HUU?**

Maelezoambayoutatupayatatusaidiakuelewajukumu la

klinikiwakatiwaujauzito.Utafitihuuunalengo la

kuchangiakatikakuboreshautoajiwahudumayaelimuyalishe bora

katikaklinikizaujauzitokatikahospitalizaummanchini Kenya.

Lengo linguine nikuhakikishwakwamba sera zilizopo, miongozonataratibuzalishe bora katikaujauzitozinawezaboreshahudumayalisheya mama katikahospitaliyaumma

**JE, KUSHIRIKI KATIKA UTAFITI HUU UTAKUGARIMU CHOCHOTE?**

Haitakugarimu pesa yoyote kushiriki katika utafiti huu

**JE , IWAPO UTAKUWA NA MASWALI YOYOTE HAPO MBELENI**

IwapoutakuwanamaswaiZaidiamawasiwasinkuhusukushirikikatikautafitihuu, tafadhalipigasimu au tumaujumbemfupikwawatafitiukitumianambariutakayopewachiniyaukurasahuu.

KwahabariZaidikuhusuhakizakokamamtafitiwa, unawezakuwasiliananamwenyekiti, kamatiyautafitiyayahospitalikuuya Kenyatta na chuo kikuu cha Nairobi. Nambariyasimu 2726300 ext 44102 email :uonknh\_erc @uonbi.ac.ke

**CHAGUO ZAKO ZINGINE NI ZIPI?**

Uamuziwakokushirikikatikautafitiniwakujirolea Unauhuruwakukataakushirikikwautafitiawawezakujiiondoakwenyeutafitiwakatiwowote

**FOMU YA IDHINI**

**TaarifayaMshiriki**

Nimesomafomuyaidhini / nimesomewanamtafiti.Nimejibiwamaswalikwalughaninayoelewa.Nimeelezwamadharanafaid ayautafitihuu.Naelewakwambakushirikikwangukwenyeutafitihuuniwakujiroleananinahakiyak ujiondoawakatiwowote.Nimechaguakushirikikatikautafitihuu.Ninaelewakwambajuhudizotezi tafanywailikuyawekamahojianohayasirinanambari fiche itatumiwakunitambua.

Kweakutiasahihikwenyefomuhii, sijatoahakizanguzakisheria.

Utakubalikushirikikatikautafiti? **NDIO/LA**

Utakubalikupeananambariyakoyasimukwamawasiliano yabaadayekuhusuutafiti: **NDIO/LA**

SahihiyaMshirikií í í í í í í í í .Tareheí í í í í í í í í .

**Taarifayamtafiti**

Nimewezakutoalmaelezo yotemuhimukuhusuutafitihuukwamshirikinanatumaikwambaameele wanaamekubalibilaushawishiwowote

Jina la mtafitií í í í í í í í í .Tareheí í í í í í í í í

Sahihiyamtafitií í í í í í í í í .Tareheí í í í í í í í í

Jukumukatikautafitií í í í í í í í í í í í í í í í í í ..